Index to data sets

Sections in which the specified data sets are analysed; bold indicates location of an introduction to the dataset and a source reference. The list is ordered by first appearance and the heading gives the data directory.

- **\Ekofisk macrofauna**
  - Ekofisk oilfield monitoring (N Sea), soft-sediment macrofauna counts & contaminant variables:
    - Sections: **1**, 4, 5, 8, 9, 10, 14

- **\Tasmania meiofauna**
  - Tasmanian soldier crab disturbance study (Australia), sandflat nematode & copepod abundance:
    - Sections: **1**, 3, 9, 10, 11

- **\WA fish diets**
  - W Australian coastal fish, dietary data, gut composition of prey categories:
    - Sections: **3**, 4, 8, 9, 17

- **\Fal benthic fauna**
  - Fal estuary mudflats (SW England), copepod, nematode & macrofauna abundance (and metal levels):
    - Sections: **4**, 10, 17

- **\N Sea biomarkers**
  - Biomarkers in flounder tissues (Southern N Sea), biochemical & histological variables:
    - Sections: **4**, 5, 9, 10, 12

- **\Denmark PSA**
  - Danish sediments, particle size distributions:
    - Sections: **4**, 9

- **\Tikus corals**
  - Tikus Island, Thousand Islands (Indonesia) over coral bleaching event, coral % area cover:
    - Sections: **5**, 16, 17

- **\Plymouth PSA**
  - Plymouth water samples, particle size distributions:
    - Section: **5**

- **\Europe groundfish**
  - Groundfish trawl surveys (NW European shelf waters), fish abundance:
    - Sections: **5**, 6, 7, 11, 16

- **\Exe nematodes**
  - Exe estuary intertidal (SW England), nematode abundance & natural environmental data:
    - Sections: **6**, 8, 9, 10, 14

- **\BC zooplankton**
  - Bristol Channel plankton net hauls (W England), zooplankton abundance:
    - Sections: **6**, 7, 8, 10

- **\Morlaix macrofauna**
  - Morlaix Bay, Amoco-Cadiz oil-spill (N coast of France), time series of macrofauna abundance:
    - Sections: **8**, 10, 14

- **\Tees macro benthos**
  - Tees Bay (NE England), time series of macrobenthic abundance in coastal sediments:
    - Sections: **8**, 9, 14

- **\World cities**
  - Great-circle distances between world cities:
    - Section: **8**

- **\Phuket corals**
  - Ko Phuket reefs under disturbance (Thailand), spatio-temporal series of coral transect cover:
    - Sections: **8**, 9, 14

- **\Messolongi diatoms**
  - Messolongi lagoon system (E Central Greece), diatom abundance & water-column environmental data:
    - Sections: 8, **13**

- **\Calafuria algae**
  - Calafuria colonisation experiments on subtidal rock (Ligurian Sea, N Italy), macroalgae cover:
    - Sections: **9**, 14

- **\Wrasse diets**

- King Wrasse diets (W Australia), gut prey composition by volume, over fish size, location and season:
  - Sections: **9**, 10

- **\NZ holdfast fauna**
  - Holdfast fauna (northern New Zealand), macrofaunal abundance at nested spatial scales:
    - Sections: **9**

- **\Frierfjord macrofauna**
  - Frierfjord/Langesundfjord (Oslofjord, Norway), subtidal soft-sediment macrofaunal abundance:
    - Sections: **10**

- **\Linnhe macrofauna**
  - Loch Linnhe pulp-mill effluent monitoring (Scotland), macrofauna abundance & biomass:
    - Sections: **10**, 16

- **\Clyde macrofauna**
  - Clyde sludge dumpground monitoring (Scotland), macrofauna abundance/biomass & contaminant variables:
    - Sections: **11**, 12, 14, 15, 16

- **\Sea-loch macrofauna**
  - Sea-lochs Loch Etive & Loch Creran (Scotland), soft-sediment macrofauna sampled over circles & transect:
    - Sections: **14**, 16

- **\Leschenault fish**
  - Leschenault estuary (W Australia), seine net samples of fish communities regionally and seasonally:
    - Section: **14**

- **\Solberg copepods**
  - Solbergstrand mesocosm, nutrient-enrichment experiments (Norway), copepod abundance:
    - Section: **14**

- **\Bermuda benthos**
  - Bermuda, Hamilton Harbour, sub-tropical macrofauna abundance:
    - Section: **15**

Revision #12
Created 12 March 2024 22:34:19 by Arden
Updated 26 February 2025 05:27:49 by Abby Miller