

Table of Contents

- **Citation**
- **Summary**
 - Introduction
 - New Statistical Methods in P8
 - New Tools & Utilities in P8
- **1. Expanded summary statistics**
 - 1.1 Expansion from P7 to P8
 - 1.2 Definitions of statistics
 - 1.3 Biotic data: summary stats
 - 1.4 Split summary stats results by groups
 - 1.5 Environmental data: summary stats
- **2. Empirical distributions**
 - 2.1 What is an empirical distribution?
 - 2.2 Example: Empirical distributions of oyster sizes
- **3. Dot plots and Violin plots**
 - 3.1 Plots of empirical densities
 - 3.2 Example: Dotplot of oyster sizes
 - 3.3 Example: Violin plot of kelp holdfast volumes
- **4. Univariate non-parametric methods**
 - 4.1 Wilcoxon signed-rank test
 - 4.2 Example: Plankton hauls
 - 4.3 Mann-Whitney U test
 - 4.4 Example: Snapper in marine reserves
 - 4.5 Kruskal-Wallis test
 - 4.6 Example: A bivalve species from Ekofisk
 - 4.7 Kolmogorov-Smirnov test
 - 4.8 Example: Sizes of oysters
 - 4.9 Test of Association
 - 4.10 Example: Ekofisk diversity
 - 4.11 Example: Associations between species
- **5. New PERMANOVA Design file**
 - Overview of new 'Design' options and tools

- **6. Allow heterogeneous dispersions in PERMANOVA**

- 6.1 Overview - Allow heterogeneity
- 6.2 ANOVA in a nutshell
- 6.3 The Behrens-Fisher problem (BFP)
- 6.4 Multivariate Behrens-Fisher problem
- 6.5 Solutions to the multivariate BFP
- 6.6 Example: one-way PERMANOVA allowing heterogeneity
- 6.7 Heterogeneity in more complex designs
- 6.8 Example: two-way crossed PERMANOVA allowing heterogeneity

- **7. Finite factors**

- 7.1 Overview - Finite factors
- 7.2 Dichotomy: fixed vs random factors
- 7.3 Not a dichotomy: a progression from fixed to random
- 7.4 Example: environmental impact on molluscs
- 7.5 Broader implications for detecting impact

- **8. Specifying Subject/Whole-plot error in PERMANOVA**

- 8.1 Designs lacking replication
- 8.2 Example: Split-plot - Woodstock vegetation
- 8.3 Example: Repeated measures - Victorian avifauna

- **9. Group covariates**

- 9.1 Why group covariables together?
- 9.2 Periodic and cyclical models
- 9.3 Example: Annual monthly cycles - B.C. macroalgae

- **10. Centroid plots**

- 10.1 Ordinations for multi-factor designs
- 10.2 Main effects plot
- 10.3 Interaction plot
- 10.4 Example: NZ fish assemblages

- **11. Residual distances**

- 11.1 What are 'residual' distances?
- 11.2 Example: Plankton (revisited)

- **12. Control charts**

- 12.1 Overview - Control charts
- 12.2 Classical univariate control chart
- 12.3 Classical multivariate control chart

- 12.4 Bivariate normal example: NZ fish
 - 12.5 Dissimilarity-based multivariate control chart
 - 12.6 Additional notes on implementing control charts
 - 12.7 Example: Birds from Grand Forks
 - **13. New standardisation options**
 - 13.1 Overview
 - 13.2 Analysing cumulative standardised data
 - 13.3 Example: Mussel sizes in the Gulf of Alaska
 - 13.4 Example: Gulf of Maine invertebrates - functional resemblance
 - **14. Create ordered groups**
 - 14.1 Overview
 - 14.2 Example: NE Pacific groundfish vs depth
 - **15. Other new tools & utilities**
 - 15.1 New default colour palette
 - 15.2 New selection options
 - 15.3 Re-name levels of a factor (or indicator)
 - 15.4 Add customised values/labels to graphical axes
 - 15.5 Split data sheet by factor/indicator
 - 15.6 Line plots for samples
 - 15.7 Output group-level stats from dispersion (or variability) weighting
 - 15.8 Output diagnostic plots from CAP
 - 15.9 New diagnostics for PCA/PCO plots
 - **16. Means plots**
 - 16.1 A plot of means with error bars
 - 16.2 Example: Fal biota (one-way case)
 - 16.3 Example: Okura macrofauna (two-way nested case)
 - 16.4 Example: Leschenault fish (two-way crossed case)
 - 16.5 Example: New Zealand snapper (three-way case)
 - **References**
-

Revision #105

Created 1 February 2026 19:58:05 by Marti

Updated 20 July 2026 02:51:56 by Marti